

Periodic Orbit Theory of the Transition to Chaos in Quantum Wells

E.² Narimanov and A. Douglas Stone

Applied Physics, Yale University, P.O. Box 208284, New Haven CT 06520-8284

(November 14, 1995; revised April 16, 1996)

An analytic theory is developed for the density of states oscillations in quantum wells in a magnetic field which is tilted with respect to the barrier planes. The main oscillations are found to come from the simplest one or two-bounce periodic orbits. We calculate their period and stability analytically and find an infinite sequence of destabilizations followed by restabilizations as the chaos parameter increases. This phenomenon explains the re-entrant frequency-doubling of the density of states peaks observed in recent magnetotunneling experiments.

PACS numbers: 05.45.+b, 72.15.Gd, 73.20.Dx

Most of our intuition about the level structure of quantum systems comes from the consideration of hamiltonians with high symmetry for which the classical motion is integrable and hence the Schrödinger equation is separable. Symmetry-breaking terms are treated by perturbation theory. This approach fails when the symmetry-breaking terms become too large and typically numerical solution is employed. An alternative approach which has been used successfully in atomic physics during the past decade is to use periodic orbit theory for the density of states (DOS) of non-integrable systems. Until recently this approach has had few applications in condensed matter physics and its power is not widely appreciated. Recently however Fromhold et al. [1] studied the tunneling I-V characteristic through a wide quantum well in a strong magnetic field oriented at an angle θ to the tunnel barriers. As they varied the voltage at fixed magnetic field ($B = 11\text{T}$) they found that resonant tunneling peaks (corresponding to the sub-band thresholds at $\theta = 0$) would abruptly change in frequency. Simple estimates show that this behavior occurs well outside the regime in which sub-band mixing can be treated by perturbation theory. Modelling of the classical dynamics in this case indicated that the motion was completely chaotic and the new structure in the tunneling density of states (DOS) was explained by the appearance of new *unstable* periodic orbits via Gutzwiller's Trace formula [2]. A very important later experiment looked at the I-V peaks in the entire (plane) parameter space of magnetic field and voltage for relatively small tilt angles [3]. At small tilt angles the frequency-doubling behavior was found at much lower magnetic field ($B \approx 5\text{T}$) but was "re-entrant", i.e. would disappear at yet higher fields (Fig.1). At higher tilt angles a more complex pattern of peak-doubling *and* tripling is observed [3]. This experiment was of particular interest because it mapped the transition regime to chaos in which stable and unstable periodic orbits exist and evolve according to the general constraints governing the transition to chaos. No previous experiments of this type exist in condensed matter

physics.

Stimulated by this experiment Shepelyansky and Stone [4] modelled its non-linear dynamics and showed that the system underwent a KAM transition to chaos described by a 2D effective map similar to the standard map [5] in which the degree of chaos was controlled by a dimensionless parameter $\beta = 2v_0B/E \sim B/\sqrt{V}$ (where $\varepsilon_0 = m^*v_0^2/2 \approx eV$ is the total injection energy). Hence at fixed tilt angle θ the degree of chaos is constant along a parabola $V \sim B^2/\beta^2$, but increases with increasing B at fixed V [6]. The initial frequency-doubling island occurs well before the dynamics becomes fully chaotic and ref. [4] noted that its onset appeared to be associated with a period-doubling bifurcation of a *stable* period-one orbit which surprisingly *restabilizes* for even higher β (chaos parameter). In this work we develop for the first time an analytic theory of the periodic orbits controlling the main DOS oscillations in the transition region to chaos. The orbits responsible for the peak-doubling at small tilt angles are *different* from those previously considered [1] and account for the re-entrant magnetotunneling behavior at small θ both qualitatively and quantitatively. At higher tilt angles these orbits remain important but others previously considered [1] make significant contributions as well.

The non-linear conductance of the well is proportional to the local DOS of the well in the vicinity of the left (emitter) barrier. Electrons tunnelling into the well at high voltages gain kinetic energy as they accelerate in the field and collide with the collector barrier. Over several collisions in the well the electron loses this energy by optic phonon emission. Therefore the tunneling resonances are substantially broadened and only are sensitive to structure in the DOS on energy scales $> \hbar/T_{\text{opt}} \sim 5\text{meV}$. The Gutzwiller Trace formula relates the contribution to the DOS which is oscillatory with energy to a sum of contributions from each periodic orbit (PO) [2]; the optic phonon broadening implies that only the few shortest PO's will give resolvable structure in the experiment. The Trace formula also implies that stable PO's typically

give larger contributions than do unstable, so that in a mainly chaotic phase space the biggest DOS oscillation comes from the few remaining stable PO's.

The origin of chaos in the tilted well is as follows. At zero tilt angle ($\theta = 0$) the acceleration along the electric field $\mathbf{E} = E\hat{z}$ and the transverse cyclotron motion decouple and are integrable. Collisions with the barriers ($v_z \rightarrow -v_z$) do not transfer energy between the cyclotron and longitudinal motion. Once the B field is tilted ($\mathbf{B} = B \cos \theta \hat{z} + \sin \theta \hat{y}$) the electron executes cyclotron motion around the $\hat{\mathbf{B}}$ direction, with a superimposed drift velocity $v_d = (E/B) \sin \theta \hat{x}$, and accelerates along $\hat{\mathbf{B}}$ due to the component $\mathbf{E} \cdot \hat{\mathbf{B}} = E \cos(\theta)$. This motion is still integrable. However now collisions with the barriers in general *do* mix the cyclotron and longitudinal energies $\varepsilon_c, \varepsilon_L$ and make the total dynamics non-integrable. The amount of energy exchange between those two degrees of freedom $\Delta\varepsilon$ depends sensitively on the *phase* of the cyclotron rotation at impact; for example if the velocity falls precisely in the $x-z$ plane $\Delta\varepsilon = 0$. Since the rate of change of the phase is $\omega_c = eB/m^*$, we find [4] increasing chaos with increasing B . Thus the restabilization of period-one orbits at higher B requires explanation.

With the choice $\mathbf{A} = (-By \cos(\theta) + Bz \sin(\theta), 0, 0)$, the system can be described by an effective Hamiltonian with two degrees of freedom [4,3]:

$$H_{eff} = \frac{p_y^2 + p_z^2}{2m} + \frac{e^2 B^2}{2m} (y \cos(\theta) - z \sin(\theta))^2 - eEz + U(z) + U(-z-d) \quad (1)$$

where $U(z < 0) = 0$, $U(z > 0) = \infty$. If $\varepsilon_0 > eV$, the electron can retain enough longitudinal energy on collision at $z = 0$ to collide again with the emitter barrier. But if $\varepsilon_i \equiv \varepsilon_0 - eV \ll eV$, (as is the case for the experiments of refs. [1,3]), then $|\Delta\varepsilon|$ is much larger at the collector barrier ($z = 0$) than at the emitter barrier. Therefore the most salient features of the classical dynamics can be described by the “1-barrier model” (1BM) [4] in which the emitter barrier $U(-z-d)$ is omitted in Eq. (1) above [7]. The periodic orbit structure of the 1BM depends only on β, θ [4]. This can be seen directly from Eq. (1): if time is measured in units of ω_c^{-1} and distance in units $E/B\omega_c$ the rescaled 1BM Hamiltonian depends only on β, θ .

For $\theta = 0$ the system is integrable and *all* the periodic orbits are of two types. First, degenerate families corresponding to $\omega_c/\omega_L = p/q$ where p, q are integers. Here $\omega_L = \pi eE/\sqrt{2m\varepsilon_L}$ is the frequency of the periodic motion in the longitudinal direction $\hat{E} = \hat{z}$ of the uniformly accelerated electron bouncing off the collector barrier. Note that this frequency *decreases* with increasing longitudinal energy as $1/\sqrt{\varepsilon_L}$ to its minimum value $\omega_L = 2\pi\omega_c/\beta$ when all the energy is longitudinal ($\varepsilon_L = \varepsilon_0$). Below this energy (determined by $\beta_{min} = p/q$) these *helical* orbits (HO) cease to exist. However there is one *isolated*

periodic orbit which exists for all energies. This *traversing orbit* (TO) has no transverse energy ($\varepsilon_L = \varepsilon_0$), its period is $T_0 = 2mv_0/eE = \beta/\omega_c$. It is distinct from all helical orbits except if $\omega_c T_0 = \beta = 2\pi p$ (p integer) when an infinitesimal increment of transverse energy generates a one-bounce HO of period T_0 . The TO determines the sub-band energy spacings of the triangular well by (integrable) semiclassical quantization, $\delta E = \hbar/T_0$ and will be relevant for the DOS oscillations at $\theta \neq 0$.

First, all one-bounce orbits for $\theta \neq 0$ must be “non-mixing”, i.e. there can be no exchange of ε_L and ε_c at the bounce as these are separately conserved between bounces and the orbit is periodic. It follows that all such orbits must collide with $v_y = 0$, since this is the condition for zero energy exchange. This geometric property of *all* the one-bounce orbits allows us to derive their periods, T , analytically. Consider such an orbit in the frame of reference moving with the drift velocity $v_d = E \sin(\theta)/B$. In this frame the orbit projected on the plane perpendicular to $\hat{B}$ is a portion of a circle with a chord removed of angular size $2\phi_0 = 2 \arcsin(v_d \omega_c T / 2v_c)$, where ϕ_0 is the cyclotron phase just after the collision, and v_c is the cyclotron velocity. Therefore, $v_c = (v_d \omega_c T / 2) \sin((2\pi - \omega_c T)/2)$. Since the motion along $\hat{B}$ between collisions is uniform acceleration, at the barrier $|\mathbf{v} \cdot \hat{\mathbf{B}}| = eE \cos(\theta) T / 2m$. Using energy conservation we obtain (for arbitrary θ)

$$\frac{(\beta/2)^2 - (\omega_c T/2)^2}{(1 - (\omega_c T/2) \cot(\omega_c T/2))^2} = \sin^2(\theta). \quad (2)$$

The solutions of Eq. (2) are shown in Fig. 2. Note that for $\beta < 2\pi$ there exists only one orbit with period $\omega_c T_0 \approx \beta$ as for the TO of the untilted system, and two more orbits appear at thresholds given by $\beta_p^t \approx 2\pi p$, $p = 1, 2, \dots \infty$. Thus we have an infinite set of *tangent bifurcations* [8] in which two new orbits (one stable and one unstable) appear; for small θ we find from Eq. (2) $\beta_p^t = 2\pi p(1 + (\theta/\pi p)^{1/3} + O(\theta^{4/3}))$. It follows from Eq. (2) that for arbitrary β there exist $2N$ (with $N = \text{integer}(\beta/\beta_N^t)$) such orbits whose periods are approximately an integer multiple of $2\pi/\omega_c$, and one orbit of longest period, T_0 , which approaches the period $\omega_c T_0 = \beta$ of the unperturbed TO. This orbit is a recognizable deformation of the TO at $\theta = 0$ (see inset, Fig. 2a), whereas the other orbits are deformations of the $\theta = 0$ helical orbits of period $\omega_c T = 2\pi p$. The asymptotes at $\omega_c T = 2\pi p$ arise because for $\theta \neq 0$ the period $T = 2\pi p/\omega_c$ is not a solution of Eq. (2) for any β . If it were a solution then in the drift frame the orbit would be a full circle, which is impossible due to the collision. The helical orbits with $\omega_c T_0 \ll \beta$ have substantial cyclotron energy and typically remain far from the emitter barrier (see inset, Fig. 2a); hence the TO is the only period-one orbit important for the tunneling DOS and it defines the “sub-band” spacing of the tilted well [3].

The stability of these periodic orbits is described by a monodromy matrix [9] M ; the orbit is unstable if $|\text{Tr}(M)| > 2$. For 1-bounce orbits we find

$$\begin{aligned} \text{Tr}(M_1) = & 4 \cos^4(\theta) (\tan^2(\theta) + (\omega_c T/2) \cot(\omega_c T/2)) \\ & \times (\tan^2(\theta) + \sin(\omega_c T)/(\omega_c T)) - 2 \end{aligned} \quad (3)$$

The results (plotted in Fig. 2) show that for small tilt angles and $\beta \leq 4\pi$ (the experimental parameter range) the TO goes unstable only in small intervals of β around $\beta = \pi, 3\pi, \dots$. This can be understood by continuity. One can easily show that at $\theta = 0$ the TO is only marginally stable ($|\text{Tr}(M)| = 2$) at $\beta = \omega_c T_0 = \pi p$, $p = 1, 2, \dots$ and is otherwise stable. Since the transition to chaos is smooth [4] with increasing θ , instability can only appear for $\theta \ll 1$ in small intervals around $\beta = \pi p$. At $\beta = \beta_p^D \approx \pi(2p-1)(1+3\theta^2/2\pi^2(p-1/2)^2)$ the instability occurs by period-doubling bifurcation and a rapid restabilization occurs at $\beta_p^R \approx \pi(2p-1)(1+\theta^2(2+1/\pi^2(p-1/2)^2))$. At $\beta \approx 2\pi p$ as noted above a stable and unstable period-one orbit appear so that *two* stable period-one orbits coexist; the one with the longer period is the TO which hence “jumps” in period. (The discontinuity in the peak data near $B \approx 10\text{T}$, $V \approx 0.4\text{V}$ in Fig. 1 occurs right at $\beta = \beta_1^t$.) After this jump the helical orbits rapidly destabilize with increasing β . For large θ or β the TO never restabilizes after its bifurcation near $\beta \approx \pi, 3\pi, \dots$, but it *always* reappears as a stable orbit at the tangent bifurcation near $\beta \approx 2\pi p$. Thus for arbitrary θ the TO undergoes an infinite sequence of destabilizations and restabilizations and stable period-one islands exist at arbitrarily high chaos parameter.

How does this explain the re-entrant behavior of the experimental DOS? As the magnetic field is increased β first passes through $\beta = \beta_1^d \approx \pi$ and the TO bifurcates: for some interval of B there exists a stable period-two orbit ($T_2 \approx 2T_0$). The Trace formula implies that if a stable period-two orbit appears there should be a large increase in the amplitude of the DOS oscillation at $\hbar/2T_0$ (see Fig. 3) giving the period-halving. Eventually the period-two orbit destabilizes by period-doubling bifurcation, its contribution to the DOS is lost, and there is a return to the original sub-band spacing determined by the TO (see Fig. 1). However in contrast to simple area-preserving 2D maps of the Chirikov type [5] the increase in chaos is not monotonic; a new stable TO appears by tangent bifurcation at yet higher B and its period-doubling bifurcation defines the next interval of peak-doubling as seen in Fig.1.

Although the 1BM explains many of the puzzling qualitative features of the data of ref. [3], for further insight and quantitative comparison we now consider the two-barrier model, Eq. (1). The PO's in the 2BM depend not only on β, θ , but also on $\gamma = \varepsilon_0/eV$. We calculate $\gamma \approx 1.1 - 1.2$ for the experiment of ref. [3] based on the standard model for the emitter state [10]. Using the

non-mixing property of the TO we can again calculate its period and stability analytically [11] and (assuming constant γ) we plot the result in Fig. 2b vs. β . Note that in the 2BM for $\theta = 0$ the TO obeys $\omega_c T_0 = \beta(1 - \sqrt{1 - 1/\gamma})$ (not β) and the behavior for small θ approaches this line except near $\beta = 2\pi p$ where the helical period-one orbits appear. These orbits have too little energy to reach the emitter barrier and hence are exactly as in the 1BM. The TO also has a very similar geometry as in the 1BM except of course for most β it collides with both barriers. However a significant difference arises in its stability properties. The order of the bifurcation and restabilization points is interchanged as a function of β (or B) (see insets, Fig. 2a,2b). This implies that the period-doubling region occurs for $\beta < \beta^d$ (smaller B) and not for $\beta > \beta^d$ as in the 1BM. This difference can be understood by noting that the period-two orbit requires substantial ε_c and hence occurs only above a threshold energy, ε_2 , determined by $T_0(\varepsilon_2, B) \approx \pi/\omega_c$. In contrast to the 1BM, in the 2BM T_0 is a *decreasing* function of energy since a higher energy collision with the emitter barrier just returns the electron to the collector sooner. Thus the high-energy region in which a stable period-two orbit exists, occurs when $\omega_c T_0 < \pi$, i.e. for smaller values of β . Denoting the point of bifurcation by β_2 , the restabilization occurs for $\beta_R < \beta_2$. At $\beta_3 < \beta_R$ the period-two orbit destabilizes and at $\beta_4 < \beta_3$ it disappears in an (inverse) tangent bifurcation with a period-two orbit which doesn't reach the emitter. Thus we identify β_4 with the onset of period-halving in the tunneling oscillations in Fig. 1, and β_2 with the return to period-one oscillations. The same sequence occurs again at higher $\beta = \beta'_2, \beta'_4$. These peak-doubling boundaries are calculated analytically from the 2BM using the estimated experimental parameters (with no fitting) and are found to agree rather well with experiment (Fig.1).

The peak behavior at higher tilt angles is more complicated, as the TO is unstable over a large interval of β and the contribution of it and its offspring to the tunneling DOS needs to be compared with other unstable orbits. Comparison to the data of ref. [3] at 38° shows peak-doubling and tripling regions in very good agreement with the bifurcation and trifurcation of the TO and a re-entrant single-peak region at high B in reasonable agreement with the restabilization of the TO after tangent bifurcation [11]. Other regions of multiple peaks appear to be associated with previously identified orbits [1] which are not bifurcations or trifurcations of the TO but whose interval of existence can be understood in a similar framework [11].

In Fig. 3 we confirm that the DOS for the tilted well does show peak-doubling in the interval predicted by our theory. Here we show a numerical calculation of the oscillatory part of the DOS appropriately scaled using Eq. (1) [11] for $\theta = 20^\circ$. As the energy increases through the classical bifurcation point the DOS oscillations clearly pick

up a secondary maximum corresponding to the period-halving, but at even higher energies the secondary maxima disappear at the predicted restabilization point for the TO. Note in Fig. 1 that the secondary maxima are not observed at lower and higher B, V where the classical theory implies a stable period-two orbit exists. However at lower fields the period $T_2 \sim 1/\omega_c$ increases, so the optic phonon damping $\sim \exp(-T_2/\tau_{opt})$ removes the contribution of the period-2 orbit from the tunneling spectra. Hence secondary maxima die out in the period-two layer at low fields (see lower curve of Fig. 3). At higher B, V we propose a different mechanism suppresses the peak-doubling. Typically, the period-two orbit has a substantial amount of cyclotron energy $\varepsilon_c \sim \varepsilon_0 - eV = (\gamma - 1)eV$, which is greater than the cyclotron energy in the emitter state ($\sim \hbar\omega_c$). Tunneling to the period-2 orbit therefore requires energy transfer from longitudinal to cyclotron motion. The amount of this energy transfer increases with increase of B and V , leading to a decrease in the tunneling matrix element for this process at high fields [11].

We thank Greg Boebinger, Harsh Mathur and Dima Shepelyansky for valuable discussions, and G.B. for access to unpublished data. This work was partially supported by NSF Grant DMR-9215065 and by a President of Russia Fellowship.




FIG. 1. Resonant tunneling peak locations for $\theta = 11^\circ$ from [3]. Pairs of solid curves enclose the regions of existence of period-two orbit for 2BM. Dashed line corresponds to the destabilization of the period-2 orbit. Variation of the effective mass with energy has been taken into account.




FIG. 2. (a) Periods of one-bounce orbits vs. $\beta = 2(v_0 B / E)$ for one-barrier model with $\theta = 11^\circ$. Solid lines denote stable orbit, dotted unstable. Dashed line is $\theta = 0$ result for TO. Triangles = period-doubling bifurcations, circles = restabilizations. Insets: (top left): Traversing orbit (TO) for $\beta = 10$ projected in y-z (left) and x-y (right) plane. (top right) same for helical orbits (HO). (Lower right): blow-up near $\beta = \pi$. (b) Same for two-barrier model with $\varepsilon/eV = \gamma = 1.1$. Lower dashed line is $\theta = 0$ result for 2BM, upper for 1BM. Grey-scale regions denote intervals of existence of period-two orbit which is born in tangent bifurcation and disappears in a backwards period-doubling bifurcation.




FIG. 3. Oscillatory contribution to DOS calculated numerically vs. action $S_0(\varepsilon_0)/\hbar$ of unperturbed TO. Note onset of secondary maxima at classical bifurcation energy (1st arrow) and disappearance at classical restabilization energy (2nd arrow). (Bottom Trace, $B = 3.53T$, $V = 0.2V$, $\theta = 20^\circ$): stable period-two interval has decreased and shifted to lower energies and secondary maxima are no longer apparent (see text). Dashed line is ratio of the period-two to period-one contributions to the DOS for $B = 5T$, $V = 0.4V$, $\theta = 20^\circ$ from Eqs. (2),(3) using the Trace formula [2]. Level-broadening is $\hbar/T_{opt} = 6.6\text{meV}$.

-
- [1] T.M.Fromhold *et al.*, Phys. Rev. Lett. **72**, 2608 (1994).
 - [2] M.C.Gutzwiller *Chaos in Classical and Quantum Mechanics* (Springer-Verlag, New York, 1990)
 - [3] G. Muller *et al.*, Phys. Rev. Lett. **75**, 2875 (1995); G.S. Boebinger *et al.*, Proc. EP2DS-IX, Surf. Sci (to be published).
 - [4] D.L.Shepelyansky and A.D.Stone, Phys. Rev. Lett. **74**, 2098 (1995)
 - [5] B.V.Chirikov, Phys. Rep. **52**, 263 (1979)
 - [6] The parabola is flattened at high voltage due to the increase of the effective mass away from the band minimum (see Fig. 1).
 - [7] With a slightly different doping profile the bottom of the well could be raised above the emitter state at zero bias allowing an experimental study of the pure 1BM.
 - [8] A.J.Lichtenberg and M.A.Lieberman, *Regular and Chaotic dynamics* (Springer-Verlag, New York, 1994)
 - [9] The stability matrix of 1-bounce periodic orbits is carried out for the effective 2D Hamiltonian (1). The full 3D orbits are marginally stable with respect to translations in x ; in the quantum DOS this leads to the familiar Landau level degeneracy which is not relevant here.
 - [10] T. Ando *et al*, Rev. Mod. Phys., **54**, 437 (1982)
 - [11] E. Narimanov and A. D. Stone, unpublished.